



Digitized by the Internet Archive
in 2007 with funding from
Microsoft Corporation

<http://www.archive.org/details/iconographyofaus58muelrich>

ICONOGRAPHY

OF

AUSTRALIAN SPECIES OF ACACIA

AND

COGNATE GENERA,

BY

BARON FERD. VON MUELLER, K.C.M.G., M. & PH.D., F.R.S.,

GOVERNMENT BOTANIST OF THE COLONY OF VICTORIA.

"SED GAUDEBITIS ET EXULTABITIS USQUE AD SEMPITERNUM IN IIS, QUÆ DEUS CREABAT."—*Prophetia Isaiae*, lxxv. 18.

FIFTH DECADE.

BY AUTHORITY:

ROBT. S. BRAIN, GOVERNMENT PRINTER, MELBOURNE.

1887.

PRICE—THREE SHILLINGS.

ICONOGRAPHY

OF

AUSTRALIAN SPECIES OF ACACIA

AND

COGNATE GENERA,

BY

BARON FERD. VON MUELLER, K.C.M.G., M. & PH.D., F.R.S.,

GOVERNMENT BOTANIST OF THE COLONY OF VICTORIA.

"SED GAUDEBITIS ET EXULTABITIS USQUE AD SEMPERNUN IN IIS, QUÆ DEUS CREABAT."—*Prophetia Isaiae*, lxxv. 18.

FIFTH DECADE.

BY AUTHORITY:

ROBT. S. BRAIN, GOVERNMENT PRINTER, MELBOURNE.

1887. 4/

Q K
495
A16
M8


v. 5

ACACIA SPATULATA.

F. v. M. in *Bentham's Flora Australiensis* ii. 356 (1864).

- 1, portion of a phyllodium.
- 2, head of unexpanded flowers.
- 3, bracts.
- 4, an unexpanded flower.
- 5, expanded flower.
- 6, front- and back-view of a stamen.
- 7, pollen-grains.
- 8, pistil.
- 9, young fruits.
- 10, lower portion of a fruit-valve with seeds.
- 11, a seed separated.
- 12, transverse section of a seed.
- 13 and 14, longitudinal sections of seeds.

All enlarged, but to various extent.




ACACIA WATTSIANA.

F. v. M. in Bentham's Flora Australiensis ii. 374 (1864).

- 1, portion of a phyllodium.
- 2, head of unexpanded flowers.
- 3, bracts.
- 4 and 5, unexpanded flowers.
- 6, expanded flower.
- 7, front- and back-view of a stamen.
- 8, pistil.
- 9, lower portion of a fruit-valve with a seed.
- 10, a seed separated.
- 11, transverse section of a seed.
- 12 and 13, longitudinal sections of seeds.

All enlarged, but to various extent.



ACACIA STRONGYLOPHYLLA.

F. v. M. fragmenta phytographiæ Australiæ viii. 226 (1874).

- 1, a phyllodium.
- 2, head of unexpanded flowers.
- 3, bracts.
- 4, unexpanded flower.
- 5, expanded flower.
- 6, front- and back-view of a stamen.
- 7, pollen-grains.
- 8, pistil.
- 9, lower portion of fruit-valve with seeds.
- 10, a seed separated.
- 11, transverse section of a seed.
- 12 and 13, longitudinal sections of seeds.

All except 1 enlarged, but to various extent.




ACACIA PYRIFOLIA.

De Candolle, Mémoire sur la famille des Légumineuses 447 (1825).

- 1, head of unexpanded flowers.
- 2, bracts.
- 3, unexpanded flower.
- 4, expanded flower.
- 5, front- and back-view of a stamen.
- 6, pollen-grains.
- 7, pistil.
- 8, lower portion of a fruit-valve with seeds.
- 9, a seed separated.
- 10, transverse section of a seed.
- 11 and 12, longitudinal sections of seeds.

All enlarged, but to various extent.



ACACIA NOTABILIS.

F. v. M. fragmenta phytographiæ Australiæ i. 6 (1858).

- 1, portion of a phyllodium.
- 2, head of unexpanded flowers.
- 3, bracts.
- 4, unexpanded flower.
- 5, expanding flower.
- 6, fully expanded flower.
- 7, front- and back-view of a stamen.
- 8, pollen-grains.
- 9, pistil.
- 10, lower portion of a fruit-valve with seeds.
- 11, a seed separated.
- 12, transverse section of a seed.
- 13 and 14, longitudinal sections of seeds.

All except 1 enlarged, but to various extent.



ACACIA LEPTOPETALA.

Bentham in Schlechtendal's *Linnæa* xxvi. 619 (1853).

- 1, portion of a phyllodium.
- 2, head of unexpanded flowers.
- 3, bracts.
- 4, unexpanded flower.
- 5, expanding flower.
- 6, fully expanded flower.
- 7, front- and back-view of a stamen.
- 8, pollen-grains.
- 9, pistil.
- 10, lower portion of a fruit-valve with seeds.
- 11, a seed separated.
- 12, transverse section of a seed.
- 13 and 14, longitudinal sections of seeds.

All enlarged, but to various extent.




ACACIA MACRADENIA.

Bentham in Mitchell's Tropical Australia 360 (1848).

- 1, portion of a phyllodium.
- 2, bracts.
- 3, expanded flower.
- 4, front- and back-view of a stamen.
- 5, pollen-grains.
- 6, pistil.
- 7, lower portion of a fruit-valve with seeds.
- 8, a seed separated.
- 9, transverse section of a seed.
- 10 and 11, longitudinal sections of seeds.

All enlarged, but to various extent.



ACACIA HARVEYI.

Bentham, Flora Australiensis ii. 368 (1864).

- 1, portion of a phyllodium.
- 2, head of unexpanded flowers.
- 3, bracts.
- 4, unexpanded flower.
- 5, expanded flower.
- 6, front- and back-view of a stamen.
- 7, pistil.
- 8, lower portion of a fruit-valve with a seed.
- 9, a seed separated.
- 10, transverse section of a seed.
- 11 and 12, longitudinal sections of seeds.

All enlarged, but to various extent.



ACACIA RETINODES.

Schlechtendal, Linnæa xx. 664 (1847).

- 1, portion of a phyllodium.
- 2, head of unexpanded flowers.
- 3, bracts.
- 4, unexpanded flower.
- 5, expanded flower.
- 6, front- and back-view of a stamen.
- 7, pollen-grain.
- 8, pistil.
- 9, lower portion of a fruit-valve with a seed.
- 10, a seed separated.
- 11, transverse section of a seed.
- 12 and 13, longitudinal sections of seeds.

All enlarged, but to various extent.




ACACIA HAKEOIDES.

A. Cunningham in Hooker's London Journal of Botany i. 354 (1842).

- 1, portion of a phyllodium.
- 2, head of unexpanded flowers.
- 3, bracts.
- 4, unexpanded flower.
- 5, expanding flower.
- 6, fully expanded flower.
- 7, front- and back-view of a stamen.
- 8, pollen-grains.
- 9, pistil.
- 10, lower portion of a fruit-valve with seeds.
- 11, a seed separated.
- 12, transverse section of a seed.
- 13 and 14, longitudinal sections of seeds.
All enlarged, but to various extent.





C O N T E N T S.

ACACIA SPATULATA.	{	ACACIA LEPTOPETALA.
WATTSIANA.		MACRADENIA.
STRONGYLOPHYLLA.		HARVEYI.
PYRIFOLIA.		RETINODES.
NOTABILIS.		HAKEOIDES.

ICONOGRAPHY

OF

AUSTRALIAN SPECIES OF ACACIA

AND

COGNATE GENERA,

BY

BARON FERD. VON MUELLER, K.C.M.G., M. & PH.D., F.R.S.,

GOVERNMENT BOTANIST OF THE COLONY OF VICTORIA.

"SED GAUDEBITIS ET EXULTABITIS USQUE AD SEMPERNUN IN HIS, QUÆ DEUB CREABAT."—*Prophetia Isaiae*, lxxv. 18.

SIXTH DECADE.

BY AUTHORITY:

ROBT. S. BRAIN, GOVERNMENT PRINTER, MELBOURNE.

1887.

PRICE—THREE SHILLINGS.

ICONOGRAPHY
OF
AUSTRALIAN SPECIES OF ACACIA
AND
COGNATE GENERA,

BY
BARON FERD. VON MUELLER, K.C.M.G., M. & PH.D., F.R.S.,
GOVERNMENT BOTANIST OF THE COLONY OF VICTORIA.

"SED GAUDEBITIS ET EXULTABITIS USQUE AD SEMPERNUM IN IIS, QUÆ DEUS CREATAT."—*Prophetia Isaiae, lxx. 18.*

SIXTH DECADE.

BY AUTHORITY:
ROBT. S. BRAIN, GOVERNMENT PRINTER, MELBOURNE.

1887. ✓

QK
495
A16
M8


v. 6

ACACIA ITEAPHYLLA.

F. v. M. in Schlechtendal's *Linnæa* xxvi. 617 (1853).

- 1, upper portion of a phyllodium.
- 2, a young flower-head with its clasping bract.
- 3, head of unexpanded flowers.
- 4, bracts.
- 5, unexpanded flower.
- 6, expanded flower.
- 7, front- and back-view of a stamen.
- 8, pollen-grain.
- 9, pistil.
- 10, lower portion of a fruit-valve with a seed.
- 11, a seed separated.
- 12, transverse section of a seed.
- 13 and 14, longitudinal sections of seeds.

All enlarged, but to various extent.



ACACIA GRAFFIANA.

F. v. M. in Melbourne Chemist, new series, ii. 118 (1887).

- 1, upper portion of a phyllodium.
- 2, young flower-head, enclosed by its bracts.
- 3, head of unexpanded flowers.
- 4, unexpanded flower.
- 5, expanded flower.
- 6, front- and back-view of a stamen.
- 7, pollen-grains.
- 8, pistil.
- 9, lower portion of a fruit-valve with seeds.
- 10, a seed separated.
- 11, transverse section of a seed.
- 12 and 13, longitudinal sections of seeds.

All enlarged, but to various extent.



ACACIA DIETRICHIANA.

F. v. M. in Melbourne Chemist, July 1882.

- 1, portion of a phyllodium.
- 2, head of unexpanded flowers.
- 3, bracts.
- 4 and 5, unexpanded flowers.
- 6, expanded flower.
- 7, front- and back-view of stamens.
- 8, pollen-grain.
- 9, pistil.
- 10, lower portion of a fruit-valve with a seed.
- 11, a seed separated.
- 12, transverse section of a seed.
- 13 and 14, longitudinal sections of seeds.

All enlarged, but to various extent.



ACACIA PRÆLONGATA.

F. v. M. in Melbourne Chemist iii. (1883).

- 1, portions of phyllodia.
- 2, head of unexpanded flowers with its bracteate stalk.
- 3, floral bracts.
- 4, unexpanded flower.
- 5, expanded flower.
- 6, front- and back-view of a stamen.
- 7, pollen-grain.
- 8, pistil.
- 9, lower portion of a fruit-valve with a seed.
- 10, a seed separated.
- 11, transverse section of a seed.
- 12 and 13, longitudinal sections of seeds.

All enlarged, but to various extent.




ACACIA STENOPHYLLA.

A. Cunningham in Hooker's London Journal i. 366 (1842).

- 1, portion of a phyllodium.
- 2, head of unexpanded flowers.
- 3, bracts.
- 4, unexpanded flower.
- 5, expanded flower.
- 6, front- and back-view of stamens.
- 7, pollen-grain.
- 8, pistil.
- 9, lower portion of a fruit-valve with a seed.
- 10, a seed separated.
- 11, transverse section of a seed.
- 12 and 13, longitudinal sections of seeds.


All enlarged, but to various extent.



ACACIA CORIACEA.

De Candolle, Mémoire sur la famille des Légumineuses 446 (1825).

- 1, portion of a phyllodium.
- 2, head of unexpanded flowers.
- 3, bracts.
- 4, unexpanded flower.
- 5, expanded flower.
- 6, front- and back-view of a stamen.
- 7, pollen-grain.
- 8, pistil.
- 9, lower portion of a fruit-valve with a seed.
- 10, a seed separated.
- 11, transverse section of a seed.
- 12 and 13, longitudinal sections of seeds.
- All enlarged, but to various extent.



ACACIA HOMALOPHYLLA.

A. Cunningham in Hooker's London Journal i. 365 (1842).

- 1, portion of a phyllodium.
- 2, head of unexpanded flowers.
- 3, bracts.
- 4, unexpanded flower.
- 5, expanded flower.
- 6, front- and back-view of stamens.
- 7, pollen-grain.
- 8, pistil.
- 9, lower portion of a fruit-valve with seeds.
- 10, a seed separated.
- 11, transverse section of a seed.
- 12 and 13, longitudinal sections of seeds.

All enlarged, but to various extent.



ACACIA PENDULA.

A.Cunningham in Loudon, hortus Britannicus 490 (1830).

- 1, portion of a phyllodium.
- 2, head of unexpanded flowers.
- 3, bracts.
- 4, unexpanded flower.
- 5, expanded flower.
- 6, front- and back-view of a stamen.
- 7, pollen-grain.
- 8, pistil.
- 9, lower portion of a fruit-valve with a seed.
- 10, a seed separated.
- 11, transverse section of a seed.
- 12 and 13, longitudinal sections of seeds.

All enlarged, but to various extent.



ACACIA HARPOPHYLLA.

F. v. M. in Bentham's Flora Australiensis ii. 389 (1864).

- 1, portion of a phyllodium.
- 2, head of unexpanded flowers.
- 3, bracts.
- 4, unexpanded flower.
- 5, expanded flower.
- 6, front- and back-view of a stamen.
- 7, pollen-grain.
- 8, pistil.
- 9, lower portion of a fruit-valve with a seed.
- 10, a seed separated.
- 11, transverse section of a seed.
- 12 and 13, longitudinal sections of seeds.
- 14, radicle, plumule and part of albumentum.

All enlarged, but to various extent.





ACACIA OSSWALDI.

F. v. M. plants of Victoria ii. 27 (1860).

- 1, portion of a phyllodium.
- 2, head of unexpanded flowers.
- 3, bracts.
- 4, unexpanded flower.
- 5, expanded flower.
- 6, front- and back-view of a stamen.
- 7, pollen-grain.
- 8, pistil.
- 9, lower portion of a fruit-valve with a seed.
- 10, a seed separated.
- 11, transverse section of a seed.
- 12 and 13, longitudinal sections of seeds.

All enlarged, but to various extent.





C O N T E N T S.

ACACIA ITEAPHYLLA.	{	ACACIA CORIACEA.
GRAFFIANA.		HOMALOPHYLLA.
DIETRICHIANA.		PENDULA.
PRÆLONGATA.		HARPOPHYLLA.
STENOPHYLLA.		OSSWALDI.

ICONOGRAPHY

OF

AUSTRALIAN SPECIES OF ACACIA

AND

COGNATE GENERA,

BY

BARON FERD. VON MUELLER, K.C.M.G., M. & PH.D., F.R.S.,

GOVERNMENT BOTANIST OF THE COLONY OF VICTORIA.

"SED GAUDEBITIS ET EXULTABITIS USQUE AD SEMPERNUM IN HIS, QUÆ DEUS CREABAT."—*Prophetia Isaiae*, lxxv. 18.

SEVENTH DECADE.

BY AUTHORITY:

ROBT. S. BRAIN, GOVERNMENT PRINTER, MELBOURNE.

1887.

PRICE—THREE SHILLINGS.

ICONOGRAPHY

OF

AUSTRALIAN SPECIES OF ACACIA

AND

COGNATE GENERA,

BY

BARON FERD. VON MUELLER, K.C.M.G., M. & PH.D., F.R.S.,

GOVERNMENT BOTANIST OF THE COLONY OF VICTORIA.

"SED GAUDESITIS ET EXULTASITIS USQUE AD SEMPERIUM IN IIS, QUÆ DEUS CREATAT."—*Prophetia Isaiae*, lxxv. 18.

SEVENTH DECADE.

BY AUTHORITY:

ROBT. S. BRAIN, GOVERNMENT PRINTER, MELBOURNE.

1887. 6

QK
495
A16
M8

v.7

ACACIA STIPULOSA.

F. v. M. in Journal of the Linnean Society iii. 119 (1858).

- 1, portion of a branchlet with phyllodia and stipules.
- 2, glandular hairlets.
- 3, head of unexpanded flowers.
- 4, bracts.
- 5, two unexpanded flowers.
- 6, expanded flower.
- 7, back- and front-view of stamens.
- 8, lower portion of fruit-valve with a seed.
- 9, a seed separated.
- 10, transverse section of a seed.
- 11 and 12, longitudinal sections of seeds.

All enlarged, but to various extent.



ACACIA LUEHMANNI.

F. v. M. fragmenta phytographiæ Australiæ xi. 116 (1881).

- 1, phyllodia.
- 2, head of unexpanded flowers.
- 3, bracts.
- 4, unexpanded flower.
- 5, expanding flower.
- 6, fully expanded flower.
- 7, front- and back-view of stamens.
- 8, pollen-grains.
- 9, pistil.
- 10, lower portion of a fruit-valve with a seed.
- 11, a seed separated.
- 12, transverse section of a seed.
- 13 and 14, longitudinal sections of seeds.

All enlarged, but to various extent.




ACACIA AMBLYGONA.

A. Cunningham in Hooker's London Journal of Botany i. 332 (1842).

- 1, portion of branchlet with stipules and a stalked head of unexpanded flowers.
- 2, phyllodia.
- 3, bracts.
- 4, unexpanded flower.
- 5, expanded flower.
- 6, front- and back-view of stamens.
- 7, pollen-grains.
- 8, pistil.
- 9, lower portion of fruit-valve with seeds.
- 10, a seed separated.
- 11, transverse section of a seed.
- 12 and 13, longitudinal sections of seeds.

All enlarged, but to various extent.



ACACIA SCALPELLIFORMIS.

Meissner in Lehmann's Plantæ Preissianæ ii. 200 (1847).

- 1, a phyllodium.
- 2, bracts.
- 3, unexpanded flower.
- 4, expanded flower.
- 5, front- and back-view of stamens.
- 6, pistil.
- 7, lower portion of fruit-valve with a seed.
- 8, a seed separated.
- 9, transverse section of a seed.
- 10 and 11, longitudinal sections of seeds.

All enlarged, but to various extent.



ACACIA PHLEBOCARPA.

F. v. M. in the Journal of the Linnean Society iii. 119 (1858).

- 1, portions of phyllodia.
- 2, head of unexpanded flowers. /
- 3, bracts.
- 4, unexpanded flower.
- 5, expanding flower.
- 6, fully expanded flower.
- 7, front- and back-view of stamens.
- 8, pollen-grain.
- 9, pistil.
- 10, lower portion of fruit-valve with seeds.
- 11, a seed separated.
- 12, transverse section of a seed.
- 13 and 14, longitudinal sections of seeds.

All enlarged, but to various extent.



ACACIA IMPRESSA.

F. v. M. in the Journal of the Linnean Society iii. 133 (1858).

- 1, portions of phyllodia.
- 2, head of unexpanded flowers.
- 3, bracts.
- 4, unexpanded flower.
- 5, expanded flower.
- 6, back- and front-view of a stamen.
- 7, pollen-grain.
- 8, pistil.
- 9, lower portion of a fruit-valve with seeds.
- 10, transverse section of a seed.
- 11 and 12, longitudinal sections of seeds.

All enlarged, but to various extent.




ACACIA FARINOSA.

Lindley in Mitchell's Three Expeditions ii. 145 (1838).

- 1, portion of a phyllodium.
- 2, head of unexpanded flowers.
- 3, bracts.
- 4, unexpanded flower.
- 5, expanded flower.
- 6, longitudinal section of an expanded flower.
- 7, front- and back-view of a stamen.
- 8, pollen-grain.
- 9, pistil.
- 10, lower portion of a fruit-valve with a seed.
- 11, a seed separated.
- 12, transverse section of a seed.
- 13 and 14, longitudinal sections of seeds.

All enlarged, but to various extent.




ACACIA SUBPOROSA.

F. v. M. Plants of Victoria ii. 24 (1860).

- 1, portion of a phyllodium.
- 2, head of unexpanded flowers.
- 3, bracts.
- 4, unexpanded flower.
- 5, expanded flower.
- 6, front- and back-view of a stamen.
- 7, pollen-grain.
- 8, pistil.
- 9, lower portion of a fruit-valve with a seed.
- 10, a seed separated.
- 11, transverse section of a seed.
- 12 and 13, longitudinal sections of seeds.

All enlarged, but to various extent.



ACACIA HETEROCLITA.

Meissner in Lehmann's Plantæ Preissianæ i. 318 (1845).

- 1, portion of phyllodia.
- 2, head of unexpanded flowers.
- 3, bracts.
- 4, unexpanded flower.
- 5, expanded flower.
- 6, back- and front-view of stamens.
- 7, pollen-grain.
- 8, pistil.
- 9, lower portion of a fruit-valve with a seed.
- 10, a seed separated.
- 11, transverse section of a seed.
- 12 and 13, longitudinal sections of seeds.
- All enlarged, but to various extent.



ACACIA SIMSII.

A. Cunningham in Hooker's London Journal of Botany i. 368 (1842).

- 1, portion of phyllodia.
- 2, head of unexpanded flowers.
- 3, bracts.
- 4, unexpanded flower.
- 5, expanding flower.
- 6, fully expanded flower.
- 7, back- and front-view of a stamen.
- 8, pollen-grains.
- 9, pistil.
- 10, lower portion of a fruit-valve with a seed.
- 11, two seeds separated.
- 12, transverse sections of seeds.
- 13 and 14, longitudinal sections of seeds.

All enlarged, but to various extent.



C O N T E N T S.

<i>ACACIA STIPULOSA.</i>	{	<i>ACACIA IMPRESSA.</i>
<i>LUEHMANNI.</i>		<i>FARINOSA.</i>
<i>AMBLYGONA.</i>		<i>SUBPOROSA.</i>
<i>SCALPELLIFORMIS.</i>		<i>HETEROCLITA.</i>
<i>PHLEBOCARPA.</i>		<i>SIMSII.</i>

ICONOGRAPHY

OF

AUSTRALIAN SPECIES OF ACACIA

AND

COGNATE GENERA,

BY

BARON FERD. VON MUELLER, K.C.M.G., M. & PH.D., F.R.S.,

GOVERNMENT BOTANIST OF THE COLONY OF VICTORIA.

"SED GAUDEBITIS ET EXULTABITIS USQUE AD SEMPERIERNUM IN IIS, QUÆ DEUS CREABAT."—*Prophetia Isaiae*, lxx. 18.

EIGHTH DECADE.

BY AUTHORITY:

ROBT. S. BRAIN, GOVERNMENT PRINTER, MELBOURNE.

1887.

PRICE—THREE SHILLINGS.

ICONOGRAPHY

OF

AUSTRALIAN SPECIES OF ACACIA

AND

COGNATE GENERA,

BY

BARON FERD. VON MUELLER, K.C.M.G., M. & PH.D., F.R.S.,

GOVERNMENT BOTANIST OF THE COLONY OF VICTORIA.

"SED GAUDEBITIS ET EXULTABITIS USQUE AD SEMPERNUM IN IIS, QUÆ DEUS CREATBAT."—*Prophetia Isaiae*, lxxv. 18.

EIGHTH DECADE.

BY AUTHORITY:

ROBT. S. BRAIN, GOVERNMENT PRINTER, MELBOURNE.

1887.

QK
495
A16
M8

v8

ACACIA ORARIA.

F. v. M. fragmenta phytographiae Australiae xi. 66 (1879).

- 1, a phyllodium.
- 2, head of unexpanded flowers.
- 3, bracts.
- 4, unexpanded flower.
- 5, expanded flower.
- 6, front- and back-view of a stamen.
- 7, pollen-grain.
- 8, pistil.
- 9, lower portion of a fruit-valve with seeds.
- 10, transverse section of a seed.
- 11, a seed separated.
- 12 and 13, longitudinal sections of seeds.

All except 1 enlarged, but to various extent.



ACACIA IMPLEXA.

Bentham in Hooker's London Journal i. 368 (1842).

- 1, portion of a phyllodium.
- 2, head of unexpanded flowers.
- 3, bracts.
- 4, unexpanded flower.
- 5, expanded flower.
- 6, front- and back-view of a stamen.
- 7, pollen-grains.
- 8, pistil.
- 9, lower portion of a fruit-valve with seed.
- 10, a seed separated.
- 11, transverse section of a seed.
- 12 and 13, longitudinal sections of seeds.

All enlarged, but to various extent.



ACACIA CYCLOPIS.

A. Cunningham in Loudon's *hortus Britannicus* 407 (1830).

- 1, portion of a phyllodium.
- 2, head of unexpanded flowers.
- 3, bracts.
- 4, unexpanded flower.
- 5, expanded flower.
- 6, front- and back-view of a stamen.
- 7, pollen-grain.
- 8, pistil.
- 9, lower portion of a fruit-valve with seed.
- 10, a seed separated.
- 11, transverse section of a seed.
- 12 and 13, longitudinal sections of seeds.

All enlarged, but to various extent.



ACACIA TRINEURA.

F. v. M. Plants of Victoria ii. 25 (1860).

- 1, portion of a phyllodium.
- 2, head of unexpanded flowers.
- 3, bracts.
- 4, unexpanded flower.
- 5, expanded flower.
- 6, front- and back-view of a stamen.
- 7, pollen-grain.
- 8, pistil.
- 9, lower portion of a fruit-valve with seed.
- 10, a seed separated.
- 11, transverse section of a seed.
- 12 and 13, longitudinal sections of seeds.

All enlarged, but to various extent.



ACACIA HOMALOCLADA.

F. v. M. fragmenta phytographiae Australiae xi. 34 (1878).

- 1, portion of a phyllodium.
- 2, head of unexpanded flowers.
- 3, bracts.
- 4, unexpanded flower.
- 5, expanded flower.
- 6, front- and back-view of a stamen.
- 7, pollen-grain.
- 8, pistil.
- 9, lower portion of a fruit-valve with seeds.
- 10, a seed separated.
- 11, transverse section of a seed.
- 12 and 13, longitudinal sections of seeds.

All enlarged, but to various extent.



ACACIA HEMIGNOSTA.

F. v. M. in Journal of the Linnean Society iii. 134 (1858).

- 1, portion of a phyllodium.
- 2, head of unexpanded flowers.
- 3, bracts.
- 4, unexpanded flower.
- 5, expanded flower.
- 6, front- and back-view of a stamen.
- 7, pollen-grain.
- 8, pistil.
- 9, lower portion of a fruit-valve with seed.
- 10, a seed separated.
- 11, transverse section of a seed.
- 12 and 13, longitudinal sections of seeds.

All enlarged, but to various extent.




ACACIA DICTYOPHLEBA.

F. v. M. fragmenta phytographiae Australiae iii. 128 (1863).

- 1, a phyllodium.
- 2, head of unexpanded flowers.
- 3, bracts.
- 4, unexpanded flower.
- 5, expanding flower.
- 6, fully expanded flower.
- 7, front- and back-view of a stamen.
- 8, pollen-grain.
- 9, pistil.
- 10, lower portion of a fruit-valve with seeds.
- 11, a seed separated.
- 12, transverse section of a seed.
- 13 and 14, longitudinal sections of seeds.

All enlarged, but to various extent.




ACACIA RETIVENEA.

F. v. M. fragmenta phytographiae Australiae iii. 128 (1863).

- 1, head of unexpanded flowers.
- 2, bracts.
- 3 and 4, unexpanded flowers.
- 5, expanded flower.
- 6, front- and back-view of a stamen.
- 7, pollen-grains.
- 8, pistil.
- 9, lower portion of a fruit-valve with seeds.
- 10, a seed separated.
- 11, transverse section of a seed.
- 12 and 13, longitudinal sections of seeds.

All enlarged, but to various extent.




ACACIA FLAVESCENS.

A. Cunningham in Hooker's London Journal i. 381 (1842).

- 1, head of unexpanded flowers.
- 2, bracts.
- 3, unexpanded flower.
- 4, expanded flower.
- 5, front- and back-view of a stamen.
- 6, pollen-grain.
- 7, pistil.
- 8, lower portion of a fruit-valve with seed.
- 9, a seed separated.
- 10, transverse section of a seed.
- 11 and 12, longitudinal sections of seeds.

All enlarged, but to various extent.



ACACIA DINEURA.

F. v. M. in Journal of the Linnean Society iii. 130 (1858).

- 1, portions of phyllodia.
- 2, head of unexpanded flowers.
- 3, bracts.
- 4, unexpanded flower.
- 5, expanded flower.
- 6, front- and back-view of a stamen.
- 7, pollen-grain.
- 8, pistil.
- 9, lower portion of a fruit-valve with seed.
- 10, a seed separated.
- 11, transverse section of a seed.
- 12 and 13, longitudinal sections of seeds.

All enlarged, but to various extent.



C O N T E N T S.

ACACIA ORARIA.	{	ACACIA HEMIGNOSTA.
IMPLEXA.		DICTYOPHLEBA.
CYCLOPIS.		RETIVENEA.
TRINEURA.		FLAVESCENS.
HOMALOCLADA.		DINEURA.

THE UNIVERSITY LIBRARY
UNIVERSITY OF CALIFORNIA, SANTA CRUZ
SCIENCE LIBRARY

This book is due on the last **DATE** stamped below.

To renew by phone, call **459-2050**.

Books not returned or renewed within 14 days
after due date are subject to billing.

SCI. LIB.

Series 2477

STORED AT NRLF

